

Phone: 732-390-7750 Fax: 844-683-2244			AsteraCancerCare.org			
PATIENT REFERRAL FORM			GENERAL			
Patient Name:			Pt. DOB:	//		
Last Patient Address:	First					
Patient City:		Pt. State:	Pt. Zi	p:		
Patient Phone: ()	-		Pt. Height:	in.		
DX:				: lbs.		
Patient Allergies:						
Insurance:			ID#:			
Referred by:			NPI#:			
Office Contact (Required):			Office Ph: ()			
			Office Fax: ()			
Office Administrator (Require	ed):	Admini	strator Ph: ()			
 Required Items/Infusion Proces Valid/signed written prescription only valid for the prescription on the prescription	ption including na 5 months, including rd es: relevant diseas tion list cific medication, a tent at your office? eded, please provid be sent to referring	g refills) se being treated r s noted on the fo de patient with a g physician.	nust be mentioned in ollowing page(s) of this Yes prescription, and have	report s form I No e patient bring on		
1. A Letter of Medical Necessity	y is required for all	l patients receivir	ng their initial infusion	at Astera (letter must		
include diagnosis, previous trea	•					
2. Benefit investigations, copa			=			
if required by the payer. <u>Right</u>						
notes providing supportive doc			•	•		
days depending on the payer ar	• •		-	•		
will update the referring doctor	-	-	-			
assistance options. For certain r company prior to rendered serv process.	-	-	-	-		
3. A pretreatment education se	ession will be provi	ided by an Advan	ced Practice Provider.			
4. Once the infusion is complet						

Page 1

Patient Name:			DOB:	_/	/	
	Last First	Middle				
	e box for medication requested, attach require Once all documentation is received, we will co <u>Required Current Lab Results</u>		-			
Note: Progress	s notes and labs must be completed within	the previous 6 months for all	new and r	enewed	l prescriptions.	
🗆 Actemra	CBC, Lipid Panel, Liver Function, PPD (pri	or to initiation)				
🗆 Benlysta (IV)) None					
🗆 Briumvi	CBC, Quantitative Serum Immunoglobuli Hep B surface antibody and Hep B core a Confirm No Vaccinations within 4 We	intibody)	erology (H	ep B su	rface antigen,	
🗆 Cimzia	CBC, Prior to initiation – PPD and Hep B Hep B core antibody)	Serology (Hep B surface antige	en, Hep B s	surface	antibody and	
Cytoxan	CBC, CMP, UA					
🗆 Entyvio	Liver Function, PPD (prior to initiation)					
Evenity	CMP, Dexa Scan within 2 years \Box C	onfirm pt. has not had an MI c	or stroke w	rithin pr	evious year	
🗆 Fasenra	Peak Flow and Other Pulmonary Function	n Tests				
🗆 Ilumya	 CBC, CMP, Prior to initiation – PPD and Hep B Serology (Hep B surface antigen, Hep B surface antibody and Hep B core antibody) Confirm up to date with vaccines and no live vaccinations within 4 weeks prior to starting therapy or have an active infection. Evaluated for malignancy. 					
□ IVIG	Hematocrit, Hemoglobin, IgG Concentra Provide dose basis in mg/kg. Doses will		-		tput	
🗆 Kisunla	Prior to initiation – confirm presence of a completed Benefits investigation and Ca		rain MRI (\	within 1	year),	
Krystexxa	G6PD Deficiency, Serum Uric Acid Levels	, Confirm Oral Urate Lowering	g Agent Dis	continu	ied	
🗆 Leqembi	Prior to initiation – confirm presence of	amyloid beta pathology and b	rain MRI (\	within 1	year)	
🗆 Leqvio	Lipid Panel					
Nucala	FEV1, Peak Flow and Other Pulmonary F	unction Tests				
	CBC, prior to initiation - Hep B Serology Hep B core antibody) Confirm No Vaccinations within 4 We		surface a	ntibody	and	
Orencia (IV)	Prior to initiation – PPD and Hep B Sero Hep B core antibody)	logy (Hep B surface antigen, F	lep B surfa	ice anti	body and	

□ Radicava None

 Remicade/Inflectra (Biosimilar might be replaced if appropriate) CBC, Liver Function, Prior to initiation – PPD and Hep B Serology (Hep B surface antigen, Hep B surface antibody and Hep B core antibody) 					
Rituxan/Ria	bni/Truxima/Ruxience (CMS approved indications only - Biosimilar might be replaced if appropriate) CBC, prior to initiation - Hep B Serology (Hep B surface antigen, Hep B surface antibody and Hep B core antibody)				
	Confirm No Vaccinations within 4 Weeks of Therapy				
Saphnelo	Up to date with all immunizations before treatment initiation and confirm no live or live attenuated vaccines are given concurrently.				
🗆 Simponi Ari	a (IV) CBC, Liver Function, Prior to initiation – PPD and Hep B Serology (Hep B surface antigen, Hep B surface antibody and Hep B core antibody)				
□ Skyrizi (IV)	Crohn's Disease Indication only - CBC, CMP (with LFTs), Prior to initiation – PPD and Hep B Serology (Hep B surface antigen, Hep B surface antibody and Hep B core antibody) Confirm No Vaccinations within 4 Weeks of Therapy or have an active infection 				
□ Soliris	Meningococcal Vaccination				
Tezspire	FEV1, Peak Flow and Other Pulmonary Function Tests				
Tysabri	MRI (MS patients), TOUCH Program Registration				
🗆 Vpriv	Gene Testing (GBA – Velaglucerase Alfa)				
🗆 Vyepti	None				
Vyvgart	Anti-AChR Antibody Positive, No Live Vaccines During Therapy				
Vyvgart Hyt	rulo (SQ – CIDP and Myasthenia Gravis) Anti-AChR Antibody Positive (Myasthenia Gravis only). No live vaccines during therapy. Confirm no active infection.				
🗆 Xolair	Asthma - Baseline Serum IgE, FEV1, Peak Flow, Other Pulmonary Function Test Chronic Idiopathic Urticaria – None				